

Sheared force-networks: anisotropies, yielding and geometry

Jacco H. Snoeijer,¹ Wouter G. Ellenbroek,² Thijs J.H. Vlugt,³ and Martin van Hecke⁴

¹*Physique et Mécanique des Milieux Hétérogènes, UMR 7636 CNRS-ESPCI,
10 rue Vauquelin, 75231 Paris Cedex 05, France*

²*Instituut-Lorentz, Universiteit Leiden, Postbus 9506, 2300 RA Leiden, The Netherlands*

³*Department of Condensed Matter and Interfaces, Debye Institute,
Utrecht University, P.O.Box 80.000, 3508 TA Utrecht, The Netherlands*

⁴*Kamerlingh Onnes Lab, Leiden University, PO box 9504, 2300 RA Leiden, The Netherlands.
(Dated: February 2, 2008)*

A scenario for yielding of granular matter is presented by considering the ensemble of force networks for a given contact network and applied shear stress τ . As τ is increased, the probability distribution of contact forces becomes highly anisotropic, the difference between average contact forces along minor and major axis grows, and the allowed networks span a shrinking subspace of all force-networks. Eventually, contacts start to break, and at the yielding shear stress, the packing becomes effectively isotropic. The size of the allowed subspace exhibits simple scaling properties, which lead to a prediction of the yield stress for packings of arbitrary contact number.

PACS numbers: 45.70.Cc, 05.40.-a, 46.65.+g

Granular media, foams and emulsions are amorphous materials which can jam and then sustain a certain amount of shear stress before yielding [1, 2, 3, 4, 5]. If one slowly increases the applied shear stress and follows the evolution of contact forces and grain locations for such systems, one encounters a rather complex set of phenomena. Firstly, before the system yields as a whole, there are non-adiabatic precursor events such as local rearrangements and microslip [6, 7, 8]. Secondly, the interparticle contact forces in these systems are organized into highly heterogeneous and fragile force networks [9, 10, 11, 12, 13] – see Fig. 1. Unraveling these shear-induced phenomena and their impact on macroscopic unjamming has remained a great challenge.

While the contact forces evolve to satisfy the applied stresses, the selection of a *specific* force network for a single numerical experiment hinges on microscopic details and packing history. On the other hand, features like contact force probabilities have turned out to be relatively insensitive to these subtle details [11, 12, 13, 14, 15], which suggests a purely statistical approach. In this Letter we characterize granular packs under shear stress, by studying *ensembles* of force networks for fixed contact networks [16, 17, 18, 19, 20, 21]. This approach is based on the fact that in jammed systems there are more contact forces than force balance equations – the ensemble simply consists of all those force networks for which the contact forces are repulsive, balance on every grain and satisfy global stress constraints, while keeping the geometry fixed. This provides a novel access to the statistics of force networks under shear stress, in which the roles of fabric and force anisotropy are separated explicitly.

We consider ensembles of sheared force networks for frictionless disks in two dimensions, for contact numbers z ranging from the lower limit $z \approx z_c = 4$ (isostatic [22]) to $z = 6$ (strongly hyperstatic). We recover a number of




FIG. 1: (a) Overview of the z - τ parameter space, where z and τ denote coordination number and shear stress respectively. For a given z , force networks cease to exist beyond a maximum τ_m (dashed line). The dots indicate the values of τ and z in panels (b-d). (b-d) Parts of the force networks, where line thicknesses represent the strength of the contact forces. These networks become highly anisotropic under shear stress.

experimental features, in particular a transition to yielding: force networks cease to exist beyond a critical yield stress τ_m . The boundary of the grey zone in Fig. 1a indicates that this maximum yield stress strongly depends on the coordination number.

In this paper we characterize the physics of sheared force networks at three levels of detail. (i) $P_\phi(f)$, the contact-angle (ϕ) resolved probability density for the contact forces f , is a natural extension of the overall force distribution $P(f)$ [11, 12, 13, 14, 15], and acts as a sensitive probe for anisotropy due to shear. (ii) The angle resolved *average* force $\bar{f}(\phi)$ is possibly the simplest characterization of shear-induced anisotropy. We find that it evolves sinusoidally as $\bar{f}(\phi) = 1 + 2\tau \sin(2\phi)$ for small τ , but that higher harmonics develop for larger shear stresses. Bounds on τ_m are obtained by requiring




FIG. 2: (a) Illustration of our geometry, showing the direction of shear stress (arrows), contact angle ϕ and major and minor axes. (b) Angle resolved force distributions for $z=5$. $P_\phi(f)$ becomes increasingly modulated with contact angle ϕ for increasing shear stress τ . (c) $P_\phi(f)$ varies qualitatively with ϕ (here $\tau=0.2$).

that $\bar{f}(\phi)$ should not become negative. (iii) The yielding curve of Fig. 1a is due to a vanishing volume of the high-dimensional space of allowed networks $\mathbf{F}_\tau$. Characterizing the size of $\mathbf{F}_\tau$ by the average distance L between random pairs of force networks, we find that L and τ are approximately related as $(L/L_m)^2 + (\tau/\tau_m)^2 = 1$. Both L_m and τ_m exhibit simple scaling laws with contact number z , $z-z_c$ and particle number N , resulting in a general prediction for the maximal shear stress τ_m that a given granular packing can sustain before yielding.

Force ensemble – The numerical results have been obtained by a recently developed ensemble technique [17, 18, 19, 20, 21]. The input of the ensemble consists of a fixed contact geometry of a 2D packing of $N = 1024$ frictionless disks of radii R_i with centers $\mathbf{r}_i$ and coordination number $z > z_c$ in a volume V , which we generated from molecular dynamics simulations of a 50:50 binary mixture of particles with size ratio 1.4 that have a purely repulsive Lennard-Jones interaction. Different densities were used to obtain different z . The system is *not* sheared and the resulting contact networks are isotropic [2, 17]. These contact networks are then kept fixed, and the positive interparticle forces between particles i and j , f_{ij} , are seen as degrees of freedom which satisfy mechanical equilibrium, restricted by the macroscopic stress $\sigma_{\alpha\beta}$:

$$\sum_j f_{ij} \frac{\mathbf{r}_i - \mathbf{r}_j}{|\mathbf{r}_i - \mathbf{r}_j|} = \mathbf{0}, \quad \sigma_{\alpha\beta} = \frac{1}{V} \sum_{\{ij\}} (f_{ij})_\alpha (\mathbf{r}_i - \mathbf{r}_j)_\beta. \quad (1)$$

In this picture there are $(zN/2)$ degrees of freedom (contact forces) constrained by $(2N+3)$ equations, leading, for $z > 4$, to an *ensemble of force networks* that form a high-dimensional convex subspace $\mathbf{F}_\tau$ [17, 18, 19]. In this paper we choose the coordinates and pressure such that




FIG. 3: Average contact force as function of contact angle ϕ and applied shear stress τ . (a) $\bar{f}(\phi)$, for $z=5$ develops a sinusoidal modulation when τ is increased. $\bar{f}(\phi)$ can be fitted well by an expression of the form $1 + 2\tau \sin(\phi) - b_2 \cos(4\phi)$; b_2 slowly increases with τ as shown in panel c. (b) $\bar{f}(\phi)$, for $z=6$ develops a strong second harmonic for large τ and approaches the limiting curve given by Eq. (4) (dashed curve). (c-d) Corresponding dependence of b_2 on τ .

$\sigma_{xx} = \sigma_{yy} = 1/2$, and consider the dimensionless shear stress $\tau = \sigma_{xy}/\sigma_{xx}$ (equivalent to the relative deviatoric stress) (see Fig. 2a).

Angle resolved force distributions – To characterize the anisotropic force networks, we introduce here the contact-angle resolved force distribution $P_\phi(f)$. This distribution is normalized such that $\int df P_\phi(f) = 1$ for all ϕ , and $1/\pi \int d\phi \int df [f P_\phi(f)] = 1$. Note that, in general, the average force for contacts in a *certain direction*, $\bar{f}(\phi) \equiv \int df [f P_\phi(f)]$, is *not* equal to one.

Figure 2b illustrates that for sheared systems $P_\phi(f)$ modulates with ϕ – this modulation has its extrema along major and minor axes. Many contact forces along the minor axis ($\phi \sim 3\pi/4$) evolve towards zero for increasing shear, effectively breaking these contacts [17] and leading to a δ -peak at $f=0$ (black area in Fig. 2b4). Fig. 2c shows examples of $P_\phi(f)$ and illustrates that for sheared systems, the qualitative *shape* of the force distribution varies with contact angle. In particular for large stresses, $P_\phi(f)$ becomes completely monotonic along the minor axis but remains peaked along the major axis. Earlier experimental and numerical findings that the total force distribution, averaged over all orientations, evolves from a peaked to a monotonically decreasing function with increasing τ are thus seen to result from averaging over strongly different $P_\phi(f)$ [1, 2, 13, 14, 15].

Analytical bounds on τ – The most basic manifestation of stress anisotropy, however, is the modulation of the average force, $\bar{f}(\phi)$, as a function of the contact orientation. This effect is clearly visible in Figs. 1, and has only recently been accessed experimentally [10, 13]. In Fig. 3 we therefore show examples of $\bar{f}(\phi)$ for various stresses and contact numbers, as obtained by the ensemble

ble. For the strongly hyperstatic case (Fig. 3b,d), it is clear that the anisotropy is limited by the requirement that $\bar{f}(\phi)$ should definitely remain positive for all ϕ . This is due to the repulsive nature of the contact forces, which requires all $f_{ij} \geq 0$. We show below that this simple criterion imposes analytical bounds on the maximum shear stress τ_m that become increasingly accurate for strongly hyperstatic packings.

This bound can be computed from the probabilistic version of Eq. (1) for isotropic contacts,

$$\sigma_{\alpha\beta} = \frac{\bar{r}N_c}{V\pi} \int_0^\pi d\phi \bar{f}(\phi) n_\alpha n_\beta, \quad (2)$$

where $\bar{r}$ is the average particle radius [23], while $(n_x, n_y) = (\cos \phi, \sin \phi)$; in the remainder we set the prefactor $\bar{r}N_c/V$ equal to unity. We Fourier expand $\bar{f}(\phi)$ as $\sum a_k \sin 2k\phi + \sum b_k \cos 2k\phi$, and anticipate that only odd k sine terms and even k cosine terms are compatible with the symmetry of a simple shear [3]. Eq. (2) then yields

$$\bar{f}(\phi) = 1 + 2\tau \sin 2\phi - b_2 \cos 4\phi + \dots, \quad (3)$$

where the coefficients of the higher order terms $b_2, \dots$ are independent of the stress tensor (Fourier modes with $k \geq 2$ yield zero upon integration). The role of the higher order terms is limited [24]: our numerics show a significant contribution to $\bar{f}(\phi)$ for large τ and z for $k = 2$ only (Fig. 3b,d). Truncating Eq. (3) at second order yields that an upper bound is reached for $\tau = \tau_m = 2/3$:

$$\bar{f}_{max}(\phi) = 1 + 4/3 \sin 2\phi - 1/3 \cos 4\phi \quad (4)$$

By extending the expression (2) to include fabric anisotropy and frictional forces, and following a similar line of reasoning, bounds can also be obtained for the completely general case [26].

Fig. 3b,d illustrate the relevance of this bound for strongly hyperstatic packings: for $z = 6$ the maximal stress is close to $2/3$, while $\bar{f}(\phi)$ approaches the limiting form Eq. (4), indicated by the dashed line. Closer to isostaticity, however, the system has much less freedom to accommodate to the shear stress, and τ_m is much smaller – as shown in Fig. 3a,c, $\bar{f}$ then does not approach zero yet. In the isostatic limit there are no adjustable degrees of freedom left, the force space $\mathbf{F}_\tau$ shrinks to a single point, and τ_m tends to zero. In general, the question is thus how τ_m depends on z (Fig. 1a).

High-dimensional ensemble – The yielding curve $\tau_m(z)$ can be determined best from the geometric properties of this high-dimensional force space. We have quantified the “size” of $\mathbf{F}_\tau$ by the Euclidian distance between random realizations (force networks), for given z and τ [19, 20]. While the distances between random points in a low dimensional space are broadly distributed, this distribution becomes increasingly sharply peaked for higher




FIG. 4: (a) The linear size L of the force space $\mathbf{F}_\tau$ as a function of shear stress τ for z as labeled (dots) are well fitted by a relation of the form $(L/L_m)^2 + (\tau/\tau_m)^2 = 1$ (solid curves). (b) Sketch of how force space $\mathbf{F}$ (represented schematically by a sphere) intersected by the constant- τ hyperplanes yields the force spaces $\mathbf{F}_{\tau_1}$ and $\mathbf{F}_{\tau_2}$ with sizes L_1 and L_2 . (c) L_m obtained by fitting Eq. (6) (circles), compared to $\sqrt{N(z-z_c)}$ (line). (d) Maximal shear stress τ_m obtained by fitting Eq. (6) (circles) is well approximated by $2(z-z_c)/z$ (line). (e) z_{eff} drops sharply when approaching τ_m .

dimensional objects as is the case here. The average distance L defined via

$$L^2(z, \tau) \equiv \langle \sum_{ij} (f_{ij} - f'_{ij})^2 \rangle, \quad (5)$$

thus serves as an effective measure for the size of $\mathbf{F}_\tau$. Here the brackets denote an average over the random pairs of force networks $\{f_{ij}\}$ and $\{f'_{ij}\}$.

In Fig. 4 we show our main findings for the main properties of the force space with τ . First, $L(\tau, z)$ can, surprisingly, be fitted by a simple relation of the form

$$(L/L_m)^2 + (\tau/\tau_m)^2 = 1, \quad (6)$$

which becomes particularly accurate for $z < 5$ (Fig. 4a). In addition, as shown in Fig. 4c-d, L_m and τ_m approach simple scaling laws:

$$\tau_m \approx 2 \frac{z - z_c}{z}, \quad (7)$$

$$L_m \approx \sqrt{N(z - z_c)}. \quad (8)$$

The scaling relations Eqs. (6-8) can be interpreted geometrically, keeping in mind that high-dimensional objects can be quite counter-intuitive (Fig. 4b). Let us consider the ensemble $\mathbf{F}$ that is obtained by applying all force balance equations and $\sigma_{xx} = \sigma_{yy} = 1/2$, but leaving σ_{xy} , and thus τ , undetermined. $\mathbf{F}$ is a convex body of dimension $D = (z - z_c)N/2 - 2$ that has $zN/2$ facets; each facet corresponds to a certain contact force being zero. One can obtain $\mathbf{F}_\tau$ from intersection of $\mathbf{F}$ with the codimension-one hyperplane given by the linear constraint $\sigma_{xy} = \tau/2$

(Eq. 1). Due to symmetry, the most “central” intersection is obtained for $\tau = 0$, while for larger values of τ , the intersection is less centered and $\mathbf{F}_\tau$ is smaller.

In fact, Eq. (8) can be recovered analytically. The scaling $L_m \propto \sqrt{D}$ with the dimension $D = N/2(z - z_c)$ of $\mathbf{F}_\tau$ is a common feature of high-dimensional convex spaces [27]. If we consider a simple approximation for $\mathbf{F}_\tau$ by ignoring the force balance equations but only requiring that all f_{ij} are positive and have $\langle f_{ij} \rangle = 1$, we obtain that $L^2 = 2D = N(z - z_c)$ [28]. Note that the relation $\tau_m \approx 2(z - z_c)/z$, implies that τ_m is proportional to the ratio between the dimension of $\mathbf{F}_\tau$ and the number of its facets $zN/2$. We have not been able to come up with a convincing argument for this scaling, however.

Finally, as τ approaches its maximum τ_m , the space $\mathbf{F}_\tau$ shrinks and $L \rightarrow 0$, so that at $\tau = \tau_m$, $\mathbf{F}_\tau$ consists of a single point. The effective contact number z_{eff} , which is defined by considering contacts broken when their force drops below a fixed small threshold [29], stays constant over most of the range of τ , but sharply drops to z_c as τ approaches τ_m (Fig. 4e).

Outlook – We have proposed an ensemble theory for force networks under shear, by exploring solutions of mechanical equilibrium for packings of fixed contact geometry. The ensemble approach provides a great conceptual simplification with respect to full numerical simulations, as it steps aside the intricate evolution of the contact network. Yet, it captures recently measured statistical properties, such as $\bar{f}(\phi)$ and the evolution of $P(f)$ [10, 13]. Furthermore, it provides an alternative description of yielding phenomena, in terms of a vanishing volume of the force phase space. Consistent with existing numerical simulations [4, 6], we found that the maximum shear stress τ_m , strongly depends on the coordination number of the packing. This dependence can be understood in terms of the geometry of the force space, and obeys a simple scaling law Eq. (7). The ensemble thus provides a new perspective for soil mechanics, in which relations between the macroscopic effective friction and micromechanical properties (density, coordination number, texture,...) play a central role. More generally, it suggests a route along which the unjamming by shear of a broad range of disordered media may be understood.

Acknowledgements We thank Alexander Morozov and Wim van Saarloos for discussion. JHS acknowledges financial support by a Marie Curie European Fellowship FP6 (MEIF-CT2003-502006), WGE from the physics foundation FOM, and TJHV and MvH from the science foundation NWO through VIDI grants.

-
- [1] A.J. Liu and S.R. Nagel, *Nature (London)* **396**, 21 (1998); V. Trappe *et al.*, *ibid.* **411**, 772 (2001).
 [2] C.S. O’Hern, S.A. Langer, A.J. Liu and S.R. Nagel, *Phys. Rev. Lett.* **86**, 111 (2001).

- [3] F. Radjai, D.E. Wolf, M. Jean and J.J. Moreau, *Phys. Rev. Lett.* **80**, 61 (1998).
 [4] F. Radjai and S. Roux (2004), in *The Physics of Granular Media*, edited by H. Hinrichsen et D. E. Wolf (Wiley-Vch, Berlin), pp. 165.
 [5] F. Alonso-Marroquín, S. Luding, H.J. Herrmann, and I. Vardoulakis, *Phys. Rev. E* **71**, 051304 (2005).
 [6] G. Combe and J.N. Roux, *Phys. Rev. Lett.* **85**, 3628 (2000).
 [7] A. Kabla *et al.*, cond-mat/0412074.
 [8] L. Silbert, private communication.
 [9] H.M. Jaeger, S.R. Nagel and R.P. Behringer, *Rev. Mod. Phys.* **68**, 1259 (1996); P.G. de Gennes, *ibid.* **71**, 374 (1999).
 [10] J. Geng, G. Reydellet, E. Clement, and R.P. Behringer, *Physica D* **182**, 274 (2003).
 [11] D.M. Mueth, H.M. Jaeger and S.R. Nagel, *Phys. Rev. E* **57**, 3164 (1998); D.L. Blair *et al.*, *ibid.* **63**, 041304 (2001).
 [12] J. Brujic *et al.*, *Faraday Discussions* **123**, 207 (2003).
 [13] T.S. Majmudar and R.P. Behringer, *Nature (London)* **435**, 1079 (2005).
 [14] E.I. Corwin, H.M. Jaeger and S.R. Nagel, *Nature (London)* **435**, 1075 (2005).
 [15] L.E. Silbert *et al.*, *Phys. Rev. E* **65**, 051307 (2002).
 [16] J.P. Bouchaud, Proceedings of the 2002 Les Houches Summer School on *Slow Relaxations and Nonequilibrium Dynamics in Condensed Matter*.
 [17] J.H. Snoeijer, T.J.H. Vlugt, M. van Hecke, and W. van Saarloos, *Phys. Rev. Lett.* **92**, 054302 (2004).
 [18] J.H. Snoeijer, T.J.H. Vlugt, W.G. Ellenbroek, M. van Hecke, and J.M.J. van Leeuwen, *Phys. Rev. E* **70**, 061306 (2004).
 [19] T. Unger, J. Kertész and D.E. Wolf, *Phys. Rev. Lett.* **94**, 178001 (2005).
 [20] S. McNamara and H.J. Herrmann, *Phys. Rev. E* **70**, 061303 (2004).
 [21] B.P. Tighe *et al.*, cond-mat/0505003
 [22] C.F. Moukarzel, *Phys. Rev. Lett.* **81**, 1634 (1998); A.V. Tkachenko and T.A. Witten, *Phys. Rev. E* **60**, 687 (1999).
 [23] Numerical simulations [3, 17] have shown that r ($\approx$ grain size) does not correlate to forces and angles, so it can be taken out of the integral.
 [24] $\bar{f}(\phi)$ is monotonic between the major and minor principal directions, and for particles with z contact forces, the width of the modulation is bounded from below by roughly $2\pi/z$ [4, 25].
 [25] H. Trodec, F. Radjai, S. Roux, and J.C. Charmet, *Phys. Rev. E* **66**, 041305 (2002).
 [26] W.G. Ellenbroek and J.H. Snoeijer, in preparation.
 [27] For example, for a hypercube of dimension D and edge 2 (such that the average value of any of its coordinates is one, to normalize similar to the forces), $L^2 = (D/4 \int_0^2 dx \int_0^2 dx' (x - x')^2) = 2D/3$.
 [28] This space is exactly that of the micro-canonical ensemble for a finite number of forces, for which $P(f) = e^{-f}$. For such a space dimension of D , we find that $L^2 = D \int_0^\infty dx \int_0^\infty dx' (x - x')^2 \exp(-x) \exp(-x') = 2D$.
 [29] Numerically, we consider a contact as broken when $f < 10^{-4} \langle f \rangle$, and we have verified that results are independent of this cutoff.